

COMPARISON OF ANALYTICAL MASS FUNCTIONS WITH NUMERICAL SIMULATIONS

Jounghun Lee^{1,2} and Sergei F. Shandarin^{1,3}

¹Department of Physics and Astronomy, University of Kansas, Lawrence, KS 66045

² Institute of Astronomy and Astrophysics, Academia Sinica, Nankang, Taipei, Taiwan

³ Theoretical Astrophysics Center, Juliane Maries Vej 30, DK-2100 Copenhagen, Denmark

taiji, sergei@kusmos.phsx.ukans.edu

Received _____; accepted _____

ABSTRACT

We present numerical testing results of our mass function derived in our previous paper, and compare the testing results with those of the popular Press-Schechter (PS) mass function. Two fiducial models are considered for the test: the scale-free power-law spectra $P(k) \propto k^n$ with spectral indices $n = -1, 0$ and the standard cold dark matter (SCDM) model with $\Omega = 1$ and $h = 0.5$. For the power-law models, we use numerical data averaged over several different output times: ten output times from two N-body realizations for the $n = -1$ power-law model; four outputs from one realization for the $n = 0$ model. While for the SCDM model, we consider four outputs separately at redshifts $z = 0, 0.43, 1.14$ and 1.86 from one large N-body simulations. The comparison results show that our mass function fits the numerical data in a much improved way over the PS one. Thus, we expect that our mass function can be a viable alternative of the PS mass function in applications to various areas.

Subject headings: cosmology: theory — large-scale structure of universe

1. INTRODUCTION

One of the most fundamental problems in cosmology is to understand the formation and evolution of the large-scale structure in the universe such as galaxies, groups and clusters of galaxies, etc. In order to understand the large-scale structure, however, it is highly desirable to have an analytical framework within which theoretical predictions for structure formation can be made. The cosmological mass function, $n(M)$ [$n(M)dM$: defined as the comoving number density of gravitationally bound structures – dark halos with mass M] provides this analytical tool since different candidate models for structure formation predict different number densities of dark halos.

Press & Schechter (1974, hereafter PS) developed for the first time an analytic formalism to evaluate the mass function. Finding a mass function requires both dynamics and statistics. Dynamically PS adopted the top-hat spherical model, according to which the collapse condition for forming dark halos is determined purely by its local average density. Statistically PS assumed that the initial density field is Gaussian, and selected dark halos from the peaks of the linear Gaussian density field.

The practical success of the PS mass function (e.g., Efstathiou et al. 1988; Lacey & Cole 1994) and the absence of viable alternatives led many authors to use it routinely in the last decade (e.g., Cole & Kaiser 1988; White & Frenk 1991; Kauffmann et al. 1993; Fan, Bahcall, & Cen 1997).

But, recently high resolution N-body simulations have shown the limitation of the PS theory: First, it has been clearly shown by several N-body simulations that the true gravitational collapse process must be nonspherical (e.g., Shandarin et al. 1995; Kuhlman et al. 1996), indicating the weakness of the PS dynamical background. Second, N-body simulations also showed that peaks of the linear density field are “poorly” correlated with the final spatial locations of dark halos (e.g., Katz, Quinn, & Gelb 1993), although the PS

formalism assumes that dark halos form in the peaks of the density field. Third, recent high-resolution numerical tests have detected that experimental results are flatter than the standard PS mass function in shape (e.g., Governato et al. 1998; Tormen 1998 and references therein ; Sheth & Tormen 1999).

Currently various attempts have been made to find a better-fitting mass function. One approach to better mass functions has been focused on finding phenomenological fitting parameters, keeping the original PS formula unchanged; for example, regarding the density threshold as a function of redshift, etc (see Governato et al. 1998; Sheth & Tormen 1999). Another approach has been on improving the dynamics of the PS formalism by implementing anisotropic collapse conditions (e.g., Monaco 1995; Audit, Teyssier, & Alimi 1997). A full analytical alternative of the PS mass function using this approach has been found by Lee & Shandarin (1998), in the frame of the Zel’dovich approximation. This approach goes along with the assumption that dark halos correspond to the third axis collapse. Indeed the mass function we found in the previous paper is both a dynamically and statistically improved version of the PS one. First, it is based on a more realistic *nonspherical* dynamical model. Second, the underlying statistical assumption that dark halos form in the local maxima of the smallest eigenvalue of the deformation tensor, λ_3 (see §2) is in general agreement with the N-body results performed by Shandarin & Klypin (1984). Third, and most importantly our mass function was shown to have desired properties like a lower peak and more high-mass halos. That is, our mass function is flatter than the PS one.

In this Letter we present numerical testing results of our mass function for the case of two fiducial models: the scale-free power-law spectra with spectral indices $n = -1, 0$, and the standard cold dark matter (SCDM) model with $\Omega = 1$ and $h = 0.5$. In §2 we briefly summarize the analytic mass function theories for the readers convenience . In §3 we

explain the N-body simulations used to produce the numerical mass functions, and compare the analytical mass functions with the numerical results. In §4 we draw a final conclusion.

2. SUMMARY OF MASS FUNCTION THEORIES

The PS theory assumes that dark halos of mass M form hierarchically in the regions where the linear Gaussian density field $\delta \equiv (\rho - \bar{\rho})/\bar{\rho}$ ($\bar{\rho}$: mean density) filtered on mass scale M reaches its threshold value δ_c for collapse:

$$n_{PS}(M) = \sqrt{\frac{2}{\pi}} \frac{\bar{\rho}}{M^2} \left| \frac{d \ln \sigma}{d \ln M} \right| \frac{\delta_c}{\sigma} \exp \left[-\frac{\delta_c^2}{2\sigma^2} \right]. \quad (1)$$

The density threshold δ_c for a flat universe is originally given by the spherical top-hat model: $\delta_c \approx 1.69$ (e.g., Peebles 1993). But in many numerical tests it has been detected that lowered δ_c (roughly 1.5) gives a better fit in the high-mass section (e.g., Efstathiou & Rees 1988; Carlberg & Couchman 1989; Klypin et al. 1995; Bond & Myers 1996). It is also worth mentioning that any PS-like formalism is least reliable in the low-mass section (Monaco 1995). This numerical detection can be understood in the following dynamical argument: Although the top-hat spherical model predicts that the gravitational collapse to “infinite” density occurs when the density reaches $\delta_c \approx 1.69$, halos in realistic case can form earlier by a rapid virialization process due to the growth of small-scale inhomogeneities (Shapiro, Iliev, & Raga 1999).

On the other hand, according to our approach (Lee & Shandarin 1998), dark halos of mass M form from the Lagrangian regions where the lowest eigenvalue λ_3 ($\lambda_3 < \lambda_2 < \lambda_1$, $\delta = \lambda_1 + \lambda_2 + \lambda_3$) of the deformation tensor d_{ij} (defined as the second derivative of the perturbation potential Ψ such that $d_{ij} = \partial^2 \Psi / \partial q_i \partial q_j$, q_i is the Lagrangian

coordinate) reaches its threshold λ_{3c} for collapse on the scale M :

$$n_{LS}(M) = \frac{25\sqrt{10}}{2\sqrt{\pi}} \frac{\bar{\rho}}{M^2} \left| \frac{d \ln \sigma}{d \ln M} \right| \frac{\lambda_{3c}}{\sigma} \left\{ \left(\frac{5\lambda_{3c}^2}{3\sigma^2} - \frac{1}{12} \right) \exp \left(- \frac{5\lambda_{3c}^2}{2\sigma^2} \right) \text{erfc} \left(\sqrt{2} \frac{\lambda_{3c}}{\sigma} \right) + \frac{\sqrt{6}}{8} \exp \left(- \frac{15\lambda_{3c}^2}{4\sigma^2} \right) \text{erfc} \left(\frac{\sqrt{3}\lambda_{3c}}{2\sigma} \right) - \frac{5\sqrt{2\pi}\lambda_{3c}}{6\pi\sigma} \exp \left(- \frac{9\lambda_{3c}^2}{2\sigma^2} \right) \right\}. \quad (2)$$

In the original derivation of our mass function, the threshold λ_{3c} for collapse has been empirically chosen to be 0.37. A similar logic used to give a dynamical explanation to the lowered δ_c of the PS formalism applies here. Although a simple extrapolation of the Zel’dovich approximation to nonlinear regime predicts that the formation of dark halos corresponding to the third axis collapse occurs at $\lambda_{3c} = 1$, the first and the second axis collapse speed up the formation of halos, which would result in a lowered λ_{3c} (see also Audit et al. 1997).

In § 3, we witness that our mass function with the original suggested value of $\lambda_3 = 0.37$ does agree with the numerical data quite well.

3. NUMERICAL vs. ANALYTICAL MASS FUNCTIONS

3.1. Comparison for Scale-Free Model

The N-body simulations of a flat matter-dominated universe for power-law spectra $P(k) \propto k^n$ with spectral indices $n = -1$ and 0 were run by White (1994) using a Particle-Particle-Particle-Mesh code with 100^3 particles in a 256^3 grid with periodic boundary conditions.

Tormen (1998, 1999) identified dark halos from the N-body simulations using a standard halo finder – the friends-of-friends algorithm with a linking length 0.2 [hereafter FOF (0.2)]. Numerical data for the $n = -1$ power-law model were obtained for 10 different output times coming from two N-body realizations, and then a final $n = -1$ numerical mass

function was obtained by taking an average over the 10 output values. While for the $n = 0$ model 4 outputs from one N-body realization were averaged to produce a final numerical mass function. For a detailed description of the simulations, see Tormen, Bouchet, & White (1997). Here we use the final average numerical mass functions for comparison data.

For the power-law spectra, the mass variance is given by the following simple form:

$$\sigma^2(M) = \left(\frac{M}{M_0} \right)^{-(n+3)/3}, \quad (3)$$

where M_0 is the characteristic mass scale¹ defined by $\sigma(M_0) = 1$. It is sometimes useful to define a filter-dependent nonlinear mass scale M_* related to M_0 by $M_* \equiv M_0(\delta_c)^{-6/(n+3)}$ for a dimensionless rescaled mass variable M/M_* such that $\sigma(M_*) = \delta_c$ (see Lacey & Cole 1994).

Figure 1 plots the fraction of mass in halos with mass M , $dF/d \ln M = (M^2/\bar{\rho})n(M)$: our mass function with $\lambda_{3c} = 0.37$ (solid line) against the averaged numerical data with Poissonian error bars, and the PS mass function with $\delta_c = 1.69$ and 1.5 (dashed and dotted lines respectively) as well. The upper panel corresponds to the $n = -1$ power-law model while the lower panel to the $n = 0$ model. As one can see, our mass function fits the numerical data much better than the PS ones for the $n = -1$ model in the high-mass section ($M > M_*$). In fact Tormen (1998) also used the spherical overdensity algorithm [SO (178)] as another halo finder, and showed that the numerical mass functions from FOF (0.2) and SO (178) are almost identical. We compared the analytical mass functions with his numerical data obtained from SO (178) and also found similar results. Whereas for the $n = 0$ model, neither of our mass function and the PS one fits the numerical data well in

¹In Lee & Shandarin (1998), the characteristic mass was notated by M_* . But here we use M_* to notate a slightly different mass scale. Readers should not be confused about this different notation of the characteristic mass.

the high-mass section. Yet in the low-mass section ($M < M_*$) our mass function fits slightly better than the PS one for this case.

3.2. Comparison for SCDM model

Governato et al. (1998) provided halo catalogs produced from one large N-body realization (comoving box size of $500h^{-1}\text{Mpc}$, 47 million particles on a 360^3 grid) of SCDM model with $\Omega = 1, h = 0.5$ for four different epochs: $z = 0, 0.43, 1.14$ and 1.86 which are respectively normalized by $\sigma(8h^{-1}\text{Mpc}) = 1.0, 0.7, 0.467$ and 0.35 . They adopted the transfer function given by Bardeen et al. (1986) and also used the FOF (0.2) halo finder. For a detailed description of the simulations, see Governato et al. (1998).

We derived the numerical mass functions from the catalogs by directly counting the number densities of halos in logarithmic scale for each epoch. In accordance with Governato et al. (1998), we consider halos more than 64 particles (corresponding to $M > 10^{14}M_\odot$) in order to avoid small-number effects of the N-body simulations. Figure 2 shows the comparison results. Our mass function with $\lambda_{3c} = 0.37$ (solid line) fits the numerical data much better than the PS ones with $\delta_c = 1.69$ and 1.5 (dashed and dotted lines respectively) for all chosen epochs.

4. CONCLUSION

We have numerically tested an analytical mass function recently derived by Lee & Shandarin (1998), and compared the results with that of the standard Press-Schechter one.

Our mass function is not just a phenomenologically obtained fitting formula but a new analytic formula derived through modification of the PS theory using a nonspherical

dynamical model. It is based on the Zel’dovich approximation taking into account the nonspherical nature of real gravitational collapse process, while the PS mass function is based on the top-hat spherical model. Consequently our mass function is characterized by the threshold value of the smallest eigenvalue of the deformation tensor, λ_{3c} while the PS one by the density threshold, δ_c .

We have shown that in the power-law model with spectral index $n = -1$ and the four different epochs of SCDM ² our mass function with $\lambda_{3c} = 0.37$ is significantly better than the PS one with $\delta_c = 1.69 - 1.5$. It fits the numerical data well especially in the high-mass section (corresponding to groups and clusters of galaxies) for these two models. Furthermore it is worth noting that in the testing results for SCDM model our mass function agrees with the data well with a consistent threshold value of 0.37 at all chosen redshifts.

On the contrary, the testing result for the $n = 0$ power-law model has shown that there are considerable discrepancies between the analytical mass functions (both of our mass function and the PS one) and the numerical data in the high-mass section. The discrepancies with theory for the $n = 0$ model, however, have been already detected (Lacey & Cole 1994). Yet in the low-mass section our mass function fits the data slightly better for this case.

Although we have tested our mass function only for two different models, given the promising testing results of our mass function demonstrated here, we conclude that it will provide a more accurate analytical tool to study structure formation. Further testings of the new mass function are obviously very desirable and will be reported in the following publications.

²At four different epochs of the SCDM model we effectively probe the dependence of the fit to the slope of the initial spectrum.

We are very grateful to Giuseppe Tormen who provided the $n = -1, 0$ numerical mass functions, and also for serving as a referee in helping to improve the original manuscript. We are also grateful to Fabio Governato’s SCDM halo catalogs and useful comments. We acknowledge the support of EPSCoR 1998 grant. S. Shandarin also acknowledges the support of GRF 99 grant from the University of Kansas and the TAC Copenhagen.

REFERENCES

- Audit, E., Teyssier, R., & Alimi, J. M. 1997, *A&A*, 325, 439
- Bardeen, J. M., Bond, J. R., Kaiser, N., & Szalay, A. S. 1986, *ApJ*, 304, 15
- Bond, J. R., & Myers, S. T. 1996, *ApJS*, 103, 1
- Carlberg, R. G., & Couchman, H. M. P. 1989, *ApJ*, 340, 47
- Cole, S., & Kaiser, N. 1988, *MNRAS*, 233, 637
- Efstathiou, G., Frenk, C. S., White, S. D. M., & Davis, M. 1988, *MNRAS*, 235, 715
- Efstathiou, G., & Rees, M. J. 1988, *MNRAS*, 230, 5p
- Fan, X., Bahcall, N. A., & Cen, R. 1997, *ApJ*, 490, L123
- Governato, F., Babul, A., Quinn, T., Tozzi, P., Bauch, C. M., Katz, N., & Lake, G. 1998, preprint (astro-ph/9810189)
- Katz, N., Quinn, T., Gelb, J. 1993, *MNRAS*, 265, 689
- kauffmann, G., White, S. D. M., & Guiderdoni, B. 1993, *MNRAS*, 264, 201
- Klypin, A., Borgani, S., Holtzman, J., & Primack, J. 1995, *ApJ*, 444, 1
- Kuhlman, B., Melott, A. L., & Shandarin, S. F. 1996, *ApJ*, 470, L41
- Lacey, C. & Cole, S. 1994, *MNRAS*, 271, 676
- Lee, J. & Shandarin, S. F. 1998, *ApJ*, 500, 14
- Monaco, P. 1995, *ApJ*, 447, 23 3
- Peebles, P. J. E. 1993, *Principles of Physical Cosmology* (Princeton: Princeton Univ. Press)

- Press, W. H., & Schechter, P. 1974, *ApJ*, 187, 425
- Shaprio, P. R., Iliev, I., & Raga, A. C. 1999, *MNRAS* in press (astro-ph/9804117)
- Shandarin, S. F., & Klypin, A. A. 1984, *Sov. Astron.*, 28, 491
- Shandarin, S. F., Melott, A. L., McDavitt, K., Pauls, J. L., & Tinker, J. 1995, *Phys. Rev. Lett.*, 75, 7
- Sheth, R. K., & Tormen, G. 1999, preprint (astro-ph/9901122)
- Tormen, G., Bouchet, F. R., & White, S. D. M. 1997, *MNRAS*, 286, 865
- Tormen, G. 1998, *MNRAS*, 290, 411
- Tormen, G. 1999, unpublished
- White, S. D. M., & Frenk, C. S. 1991, *ApJ*, 379, 25
- White, S. D. M. 1994, unpublished




Fig. 1.— The square dots represent the numerical mass function with poissonian error bars. The solid line is the LS mass function with $\lambda_{3c} = 0.37$ while the dashed, the dotted lines are the PS mass functions with $\delta_c = 1.69, 1.5$ respectively. The upper and the lower panels correspond to the $n = -1$ and the $n = 0$ power-law models respectively. See also the top left panel of Fig.2 in Tormen (1998).




Fig. 2.— The square dots represent the numerical data for the case of SCDM model with $\Omega = 1, h = 0.5$. The solid line is our mass function with $\lambda_{3c} = 0.37$, and the dashed, the dotted lines are the PS mass functions with $\delta_c = 1.69, 1.5$ respectively.